

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C07K 16/12</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/11037</b> <b>(43) International Publication Date:</b> 2 March 2000 (02.03.00)
<b>(21) International Application Number:</b> PCT/US99/19034 <b>(22) International Filing Date:</b> 20 August 1999 (20.08.99)  <b>(30) Priority Data:</b> 60/097,470 21 August 1998 (21.08.98) US 60/102,179 28 September 1998 (28.09.98) US  <b>(71) Applicant (for all designated States except US):</b> THE REGENTS OF THE UNIVERSITY OF CALIFORNIA [US/US]; 1111 Franklin Street, 12th floor, Oakland, CA 94607-5200 (US).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> SHI, Wenyan [CN/US]; 7011 Kenwood Avenue, Los Angeles, CA 90045 (US). HUME, Wyatt [AU/US]; 2808 Nicada Avenue, Los Angeles, CA 90077 (US).  <b>(74) Agent:</b> BERMAN, Charles; Oppenheimer Wolff & Donnelly LLP, 2029 Century Park East, Suite 3800, Los Angeles, CA 90067-3024 (US).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> MONOCLONAL ANTIBODIES SPECIFIC FOR STREPTOCOCCUS MUTANS, AND USES THEREOF  <b>(57) Abstract</b>  The invention describes three monoclonal IgG antibodies, referred to as SWLA1, SWLA2, and SWLA3, which appear to recognize a species-specific lipooligosaccharide or lipopolysaccharide on the cell surface of <i>S.mutans</i> . The invention also describes a rapid method of detection of <i>S.mutans</i> without the need for prior growth of the bacteria in culture. The invention further describes method of utilizing these antibodies for rapidly quantitatively detecting <i>S.mutans</i> . These methods are sensitive enough to detect the presence of a single <i>S.mutans</i> bacterial cell. These methods can be widely used in the clinical diagnosis and treatment of dental caries in humans.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## MONOCLONAL ANTIBODIES SPECIFIC FOR STREPTOCOCCUS MUTANS, AND USES THEREOF

by

5

Wenyuan Shi &amp; Wyatt Rory Hume

CROSS-REFERENCES

10

This application claims priority from a United States provisional application, United States Application Serial No. 60/102,179, filed September 28, 1998.

BACKGROUND OF THE INVENTION

15

The present invention relates to novel antibodies to the *Streptococcus mutans* bacteria that are naturally found in the mouth, and play a role in the development of dental caries. The invention relates to methods of detection of *S. mutans* using the antibodies of the invention or fragments or derivatives thereof. The invention also relates to diagnosing, monitoring, treating and protecting the teeth from dental caries using the antibodies of the invention or fragments or derivatives thereof.

20

Throughout this application, various publications are referenced within parentheses and cited at the end of the application. The disclosures of these publications are hereby incorporated by reference herein in their entireties.

25

Currently human dental caries, or cavities, are detected by changes in translucency, color, hardness or X-ray density of teeth. These technologies have limitations both in specificity and reproducibility. Further, they do not show, at a single time point, whether or not the disease is active.

30

-2-

The bacterium *Streptococcus mutans*, or *S. mutans* (named and described by Clark in 1924) is known to be a prime etiologic agent for the initiation and progression of human dental caries. (Fitzgerald and Keyes, 1960; Loesche, 1982; Loesche, 1986; Tanzer, 1997). *S. mutans* is one of the primary factors in acid  
5 dissolution of the apatite (mineral) component of the enamel then the dentin, or of the cementum then the dentin. A strong correlation between the proportion of *S. mutans* in dental plaque or in saliva relative to other bacterial species and the presence or risk of future outbreaks of dental caries has been documented, (Tanzer, 1997; Krasse, 1988). Therefore, *S. mutans* in plaque or saliva can serve as an index for both caries activity  
10 state and caries risk or susceptibility (Loesche et al., 1975; Ellen, 1976; Krasse, 1985; Krasse, 1988). These indices play an increasingly important role in the diagnosis and treatment of dental caries (Hume, 1993; Mundorff et al., 1993; Van Houte, 1993).

Present techniques of detection and quantitative determination of *S. mutans* include bacterial culture with selective media using either broth or agar plate  
15 systems (Ellen, 1976; Loesche, 1982) and polymerase chain reaction techniques (Igarashi et al., 1996). Each of these methods requires significant time (on the order of days), well trained personnel and sophisticated equipment to perform. Consequently, existing techniques are relatively expensive and time consuming.

Alternatively, monoclonal antibody based detection methods allow a  
20 rapid and accurate, yet economic quantitative measurement of the presence of bacterial cells, and have significant advantages compared to traditional culture sensitivity assays or polymerase chain reaction (PCR) techniques. Bacteria produce unique polysaccharide structures of lipooligosaccharide or lipopolysaccharide or other polysaccharides, among a large range of other chemical components and products, on  
25 their cell surfaces. Monoclonal antibodies can be raised against these chemical structures, using standard hybridoma techniques (Kohler and Milstein, 1975). The sensitivity and accuracy of this method is largely dependent on the specificity of the monoclonal antibody produced. By making and screening large numbers of hybrid cell lines, one can find certain monoclonal antibodies which are species-specific for the

-3-

desired bacteria, i.e. the monoclonal antibodies ICL 11 and ICL12 recognize the 0139 antigen of *Vibrio cholerae* (Hasan, 1994)). These monoclonal antibodies can be linked to various detection systems including, for example, fluorescent reagents, colorimetric reagents or coagglutination reagents. The resulting labeled antibodies can specifically  
5 bind to the desired bacterium in any sample, and rapidly present the result through the linked detection systems (Harlow and Lane, 1988).

TABLE 1

	PCR	Culture Sensitivity	monoclonal antibody
Reporting time	2 days send out samples	7-10 days send out samples	3-10 min instant results at chairside
Cost	~\$100	~\$100	less than \$10
Precision	Species level	genus level	Species level
Test sensitivity	few hundred cells regardless of viability	only cultivable cells need viable cells	few hundred cells regardless of viability
Accuracy	>90%	~50%	>90%
Lab requirement	yes	yes	no

Due to overwhelming advantages, monoclonal antibody-based detection methods have been widely used in medical microbiology. At present, there are nearly one hundred different monoclonal antibody based detection methods available to  
5 diagnose various pathogenic bacteria. However, to date there have been no such methods available for the detection of dental caries. While other investigators have made monoclonal antibodies to *S. mutans*, these monoclonal antibodies are not suited for diagnostic or clinical uses because they are not species specific for *S. mutans* and cannot detect the presence of *S. mutans* at very low levels. Almost all previous  
10 monoclonal antibodies were made against surface proteins of *S. mutans* (i.e. glucosyltransferase, agglutinin, surface antigen PI, etc.). Similar proteins can also be found on other Streptococci species. Thus, these monoclonal antibodies can not be used to distinguish *S. mutans* from other Streptococci species. To diagnose dental caries it is important to be able to discern the *S. mutans* strain from other bacteria that  
15 may be present. One needs to find a monoclonal antibody that is specific for a unique epitope specific to the *S. mutans* cell surface.

### SUMMARY OF THE INVENTION

20

Accordingly, the invention describes three monoclonal IgG antibodies, referred to as SWLA1, SWLA2, and SWLA3, which appear to recognize a species-specific polysaccharide on the cell surface of *S. mutans*. The invention also describes a rapid method of detection of *S. mutans* without the need for prior growth of bacteria in  
25 culture. The invention further describes methods of utilizing these antibodies for rapidly quantitatively detecting *S. mutans*. These methods are sensitive enough to detect the presence of a single *S. mutans* bacteria cell. These methods can be widely used in the clinical diagnosis and treatment of dental caries. Yet another embodiment of the invention is a diagnostic kit containing the monoclonal antibodies of the  
30 invention and reagents for detecting binding of the antibodies to *S. mutans* cells on teeth in or in a sample from a subject.

In particular, one aspect of the present invention is a monoclonal antibody that specifically binds an antigen on the surface of *Streptococcus mutans* and which is the monoclonal antibody produced by a hybridoma deposited with the American Type Culture Collection as ATCC No. HB12559, and which is designated SWLA1.

Another aspect of the present invention is a monoclonal antibody that specifically binds an antigen on the surface of *S. mutans* and which is the monoclonal antibody produced by a hybridoma deposited with the American Type Culture Collection as ATCC No. HB 12560, and which is designated SWLA2.

Yet another aspect of the present invention is a monoclonal antibody that specifically binds an antigen on the surface of *S. mutans* and which is the monoclonal antibody produced by a hybridoma deposited with the American Type Culture Collection as ATCC No. HB 12258, and which is designated SWLA3.

### BRIEF DESCRIPTION OF THE DRAWINGS

The following invention will become better understood with reference to the specification, appended claims, and accompanying drawings, where:

Figure 1 shows the detection of *S. mutans* SWLA1 with fluorescence microscopy; the *S. mutans* cells (ATCC25175) were labeled as described in Example 1, *infra*; 1(a) shows a regular phase-contrast light microscope image of a bacteria mixture containing *S. rattus*, *S. gordonii*, *S. mitis*, *S. sanguis*, *E. coli*, and *S. mutans* with FITC conjugated monoclonal antibodies; 1(b) shows the same group of bacteria under fluorescent lighting; the bright spots are all *S. mutans*, indicating that the SWLA1 monoclonal antibody specifically recognizes *S. mutans*.

Figure 2 shows flow cytometry analysis of *S. mutans* SWLA1, as described in example 1, *infra*; 2(a) shows *S. mutans* (ATCC25175) without SWLA1 and



FICT-linked goat-anti-mouse IgG antibody; 2(b) shows *S. mutans* with SWLAI and FICT-linked goat-anti-mouse IgG antibody; 2(c) shows *F. nucleatum* (ATCC25586) without SWLAI and FICT-linked goat-anti-mouse IgG antibody; 2(d) shows *F. nucleatum* with SWLAI and FICT-linked goat-anti-mouse IgG antibody; 2(e) shows *T. denticola* (ATCC33520) without SWLAI and FICT-linked goat-antibody mouse IgG antibody; 2(f) shows *T. denticola* with SWLAI and FICT-linked goat-anti-mouse IgG antibody; F is the area containing bacterial particles labeled with FITC-linked monoclonal antibody.

10

## DETAILED DESCRIPTION OF THE INVENTION

The invention describes three species-specific monoclonal IgG antibodies, referred to as SWLA1, SWLA2, and SWLA3, which recognize a species-specific epitope on the cell surface of *S. mutans*, and conjugates thereof, which appears to be a species-specific polysaccharide. The invention includes methods of using the monoclonal antibodies to detect quantity and presence of *S. mutans* to monitor the onset and severity of dental caries.

The monoclonal antibodies SWLA1, SWLA2, and SWLA3 can be prepared by hybridoma fusion techniques or by techniques that utilize EBV-immortalization technologies. Hybridoma fusion techniques were first introduced by Kohler and Milstein (see, Kohler and Milstein, (1975); Brown et al., (1981); Brown et al., (1980); Yeh et al., (1976); and Yeh et al., (1982)).

25

These techniques involve the injection of an immunogen (e.g., purified antigen or cells or cellular extracts carrying the antigen) into an animal (e.g., a mouse) so as to elicit a desired immune response (i.e., production of antibodies) in that animal. For example, *S. mutans* may be used as the immunogen. In the illustrative example herein, *S. Mutans* strain ATCC25175 was used as the immunogen. The cells are injected repeatedly, for example, into a mouse and, after a sufficient time, the mouse is

30

sacrificed and somatic antibody-producing cells are obtained. The use of other mammalian models, for example rat, rabbit and frog somatic cells, is also possible. The cell chromosomes encoding desired immunoglobulins are immortalized by fusing them with myeloma cells, generally in the presence of a fusing agent such as polyethylene glycol (PEG). Any of a number of myeloma cell lines may be used as a fusion partner according to standard techniques; for example, the P3-NSI/1-Ag4-1, P3-x63-Ag8.653 or Sp2/0-Ag14 myeloma lines. These myeloma lines are available from the American Type Culture Collection (ATCC), in Rockville, Maryland.

The resulting cells, which include the desired hybridomas, are then grown in a selective medium, such as HAT medium, in which unfused parental myeloma or lymphocyte cells eventually die. Only the hybridoma cells survive and can be grown under limiting dilution conditions to obtain isolated clones. The supernatants of the hybridomas are screened for the presence of antibody of the desired specificity, e.g., by immunoassay techniques using the antigen that has been used for immunization. Positive clones can then be subcloned under limiting dilution conditions and the monoclonal antibody produced can be isolated. Various conventional methods exist for isolation and purification of the monoclonal antibodies so as to free them from other proteins and other contaminants. Commonly used methods for purifying monoclonal antibodies include ammonium sulfate precipitation, ion exchange chromatography, and affinity chromatography (Zola et al. (1982)). Hybridomas produced according to these methods can be propagated *in vitro* or *in vivo* (in ascites fluid) using techniques known in the art (see, generally, Fink et al., supra, 1984).

Generally, the individual cell line may be propagated *in vitro*, for example in laboratory culture vessels, and the culture medium containing high concentrations of a single specific monoclonal antibody can be harvested by decantation, filtration or centrifugation. Alternatively, the yield of monoclonal antibody can be enhanced by injecting a sample of the hybridoma into a histocompatible animal of the type used to provide the somatic and myeloma cells for the original fusion. Tumors secreting the specific monoclonal antibody produced by

the fused cell hybrid develop in the injected animal. The body fluids of the animal, such as ascites fluid or serum, provide monoclonal antibodies in high concentrations. When human hybridomas or EBV-hybridomas are used, it is necessary to avoid rejection of the xenograft injected into animals such as mice. Immunodeficient or nude mice may be used or the hybridoma may be passaged first into irradiated nude mice as a solid subcutaneous tumor, cultured in vitro and then injected intraperitoneally into pristane primed, irradiated nude mice which develop ascites tumors secreting large amounts of specific human monoclonal antibodies.

For certain therapeutic applications chimeric (mouse-human) or human monoclonal antibodies may be preferable to murine antibodies, because patients treated with mouse antibodies generate human antimouse antibodies. (Shawler et al., (1985)). Chimeric mouse-human monoclonal antibodies reactive with *S. mutans* can be produced, for example, by techniques developed for the production of chimeric antibodies (Oi et al., (1986); Liu et al., (1987)). Accordingly, genes coding for the constant regions of the SWLA1, SWLA2, or SWLA3 antibody molecule are substituted with human genes coding for the constant regions of an antibody with appropriate biological activity (such as the ability to selectively bind *S. mutans*).

Novel antibodies of mouse or human origin can be also made that are analogous to the SWLA1, SWLA2, or SWLA3 antibody and that have the appropriate biological functions. These antibodies can have complementarity-determining regions (CDRs) that are identical to one of SWLA1, SWLA2, or SWLA3. Alternatively, these antibodies can bind an antigen on the surface of *S. mutans* and can compete at least about 80% as effectively on a molar basis with at least one of SWLA1, SWLA2, or SWLA3 as SWLA1, SWLA2, or SWLA3 for binding to the antigen on the surface of *S. mutans*. These antibodies have substantially no reactivity with any of the following bacterial strains: *Streptococcus rattus* ATCC19645, *Streptococcus gordonii* ATCC10558, *Streptococcus gordonii* ATCC13396, *Streptococcus mitis* ATCC49456, *Streptococcus sobrinus* ATCC33478, *Streptococcus sobrinus* 6715, *Streptococcus sanguis* ATCC10556, *Streptococcus sanguis* ATCC49295, *Streptococcus anginosus*

ATCC33397, *Lactobacillus acidophilus* ATCC4356, *Lactobacillus casei* ATCC4646, *Actinobacillus actinomycetemcomitans* ATCC 33384, *Porphyromonas gingivalis* ATCC33277, *Prevotella intermedia* ATCC49046, *Bacteroides forsythus* ATCC43047, *Eikenella corrodens* ATCC23834, *Fusobacterium nucleatum* ATCC25586, *Treponema*  
5 *denticola* ATCC33520, *Campylobacter rectus* ATCC33238, *Myxococcus xanthus* DZ2, and *Escherichia coli* HB101. Preferably, the monoclonal antibody competes at least about 90% as effectively on a molar basis.

For example, human monoclonal antibodies may be made by using the  
10 antigen, e.g. the portion of the polysaccharide on the cell surface of *S. mutans*, which binds the antibodies SWLA1, SWLA2, or SWLA3 of the invention, to sensitize human cells to the antigen in vitro followed by EBV-transformation or hybridization of the antigen-sensitized cells with mouse or human cells, as described by Borrebaeck et al. (1988).

15

The antibodies of this invention were produced via hybridoma techniques. NSL/Ag4.1 mouse myeloma cell line was used as a fusion partner, and whole cells of type c *S. mutans* strain ATCC25175 were used as the immunogen as described in the Example, *infra*. The hybridomas produced were screened. When this  
20 was performed three species-specific monoclonal antibodies against *S. mutans* were obtained designated SWLA1, SWLA2, and SWLA3. The hybridomas, producing the SWLA1, SWLA2, and SWLA3 antibodies, have been deposited with the ATCC, Rockville, Maryland, and are identified as follows:

25	SWLA1	Accession No.: HB-12559
	SWLA2	Accession No.: HB-12560
	SWLA3	Accession No.: HB-12558

The three monoclonal antibodies were found to be of the IgG subtype.  
30 Western blot and other biochemical analysis shows that they are against unique

polysaccharide on cell surface of *S. mutans*. Western blot techniques are well known by one skilled in the art (Golub, E. S. and D. R. Green, 1991).

These monoclonal antibodies are very different from the monoclonal  
5 antibodies found in the prior art. Those antibodies were made against surface proteins that are found on many other bacteria and not the species-specific polysaccharide, which the instant antibodies recognize. This difference makes the prior antibodies acceptable for research purposes but not diagnostic purposes (Carlen and Olsson, 1995; Chia et al., 1993; Fukushima et al., 1993; Brady, et al., 1991). Due to the specificity for  
10 *S. mutans*, the monoclonal antibodies of this invention are particularly useful for diagnosis and treatment of human dental caries. Because the monoclonal antibody of the invention are able to detect low numbers of *S. mutans* cells in small samples they are able to be used to screening for *S. mutans* cells. These monoclonal antibodies also permit the development of simple and inexpensive dental caries detection methods that  
15 could be used for caries risk assessment at a dentist's chairside or in the patient's household.

The most preferred antibodies will selectively bind to *S. mutans* and will not bind (or will bind weakly) to non- *S. mutans* bacteria. The antibodies that are  
20 particularly contemplated include monoclonal antibodies as well as fragments of monoclonal antibodies containing an *S. mutans* antigen-binding domain. The invention also encompasses antibody fragments that specifically recognize *S. mutans*. As used herein, an antibody fragment is defined as at least a portion of the immunoglobulin molecule which binds to its target, i.e., the antigen binding region on the *S. mutans*.  
25 This includes Fv, Fab, Fab' and F(ab)'<sub>2</sub> fragments of appropriate specificity.

The invention further includes a human monoclonal antibody that specifically binds an antigen found on the surface of *S. mutans*. The antigen that is bound is one of those bound by at least one of the monoclonal antibody produced by a  
30 hybridoma deposited with the American Type Culture Collection as ATCC No. HB 12559 and designated SWLA1, the monoclonal antibody produced by a hybridoma

deposited with the American Type Culture Collection as ATCC No. HB 12560 and designated SWLA2, and the monoclonal antibody produced by a hybridoma deposited with the American Type Culture Collection as ATCC No. HB 12558 and designated SWLA3. The human monoclonal antibody has substantially no reactivity with any of the following bacterial strains: *Streptococcus rattus* ATCC19645, *Streptococcus gordonii* ATCC10558, *Streptococcus gordonii* ATCC13396, *Streptococcus mitis* ATCC49456, *Streptococcus sobrinus* ATCC33478, *Streptococcus sobrinus* 6715, *Streptococcus sanguis* ATCC10556, *Streptococcus sanguis* ATCC49295, *Streptococcus anginosus* ATCC33397, *Lactobacillus acidophilus* ATCC4356, *Lactobacillus casei* ATCC4646, *Actinobacillus actinomycetemcomitans* ATCC 33384, *Porphyromonas gingivalis* ATCC33277, *Prevotella intermedia* ATCC49046, *Bacteroides forsythus* ATCC43047, *Eikenella corrodens* ATCC23834, *Fusobacterium nucleatum* ATCC25586, *Treponema denticola* ATCC33520, *Campylobacter rectus* ATCC33238, *Myxococcus xanthus* DZ2, and *Escherichia coli* HB101.

15

Methods for the preparation of human monoclonal antibodies are known in the art and include phage display techniques and isolation of human hybridomas using B lymphocytes from patients producing antibodies against *S. mutans*, as well as *in vitro* immunization techniques. Such techniques are well known in the art and are described, for example, in C.A.K. Borrebaeck, ed., "Antibody Engineering" (2d ed., Oxford University Press, New York, 1995), incorporated herein by this reference.

20

The invention further includes chimeric antibodies, including humanized antibodies. This includes chimeric antibodies that have complementarity-determining regions that are identical with the complementarity-determining regions of one of:

25

(a) a monoclonal antibody produced by a hybridoma deposited with the American Type Culture Collection as ATCC No. HB 12559 and designated SWLA1;

30

(b) a monoclonal antibody produced by a hybridoma deposited with the American Type Culture Collection as ATCC No. HB 12560 and designated SWLA2; or

(c) a monoclonal antibody produced by a hybridoma deposited with the American Type Culture Collection as ATCC No. HB 12558 and designated SWLA3.

5 Also within the scope of the invention are chimeric antibodies that have complementarity-determining regions that are identical with the complementarity-determining regions of an antibody that binds an antigen on the surface of *S. mutans* and can compete at least about 80% as effectively on a molar basis with at least one of SWLA1, SWLA2, or SWLA3 as SWLA1, SWLA2, or SWLA3 for binding to the  
10 antigen on the surface of *S. mutans*. These chimeric antibodies, as described above, have substantially no reactivity with any of the following bacterial strains:  
*Streptococcus rattus* ATCC19645, *Streptococcus gordonii* ATCC10558, *Streptococcus gordonii* ATCC13396, *Streptococcus mitis* ATCC49456, *Streptococcus sobrinus* ATCC33478, *Streptococcus sobrinus* 6715, *Streptococcus sanguis* ATCC10556,  
15 *Streptococcus sanguis* ATCC49295, *Streptococcus anginosus* ATCC33397, *Lactobacillus acidophilus* ATCC4356, *Lactobacillus casei* ATCC4646, *Actinobacillus actinomycetemcomitans* ATCC 33384, *Porphyromonas gingivalis* ATCC33277, *Prevotella intermedia* ATCC49046, *Bacteroides forsythus* ATCC43047, *Eikenella corrodens* ATCC23834, *Fusobacterium nucleatum* ATCC25586, *Treponema denticola*  
20 ATCC33520, *Campylobacter rectus* ATCC33238, *Myxococcus xanthus* DZ2, and *Escherichia coli* HB101.

These chimeric antibodies specifically bind an antigen on the surface of *S. mutans* and which have at least a portion of the amino acid sequence of the heavy  
25 chain or the light chain of a different species origin than the species origin of the complementarity-determining regions. In one alternative, at least a portion of the amino acid sequence of the heavy chain or the light chain is of human origin so that the chimeric antibody is a humanized antibody. In a version of this alternative, substantially all of the amino acid sequences of the heavy chain and the light chain  
30 outside the complementarity-determining regions are of human origin.

As indicated, chimeric antibodies according to the present invention may have a non-human antigen-binding site and a humanized effector binding region. The non-human antigen-binding portion may include, but is not limited to, a murine, canine, feline or other veterinary model or other mammalian antigen-binding site.

5

Methods for producing chimeric antibodies, including humanized antibodies, are well known in the art and are described, for example, in C.A.K. Borrebaeck, ed., "Antibody Engineering" (2d ed., Oxford University Press, New York, 1995), incorporated herein by this reference.

10

The invention further includes single-chain binding fragments, known generally as sFv, that have the appropriate specificity for the antigen on the cell surface of *S. mutans* as defined above. Methods for preparing such sFv are generally known in the art and are described, for example, in C.A.K. Borrebaeck, ed., "Antibody Engineering" (2d ed., Oxford University Press, New York, 1995), incorporated herein by this reference.

15

The specificity of the SWLA1, SWLA2, and SWLA3 antibodies for *S. mutans* antigen make these antibodies excellent markers for screening, diagnosis, prognosis, and follow-up assays, imaging methodologies, and therapeutic methods in the management of dental caries. The invention provides various immunological assays useful for the detection of *S. mutans* and for the diagnosis of dental caries or the risk thereof. This includes various immunological assay formats well known in the art, including, but not limited to, various types of radioimmunoassays, enzyme-linked immunosorbent assays (ELISA), enzymelinked immunofluorescent assays (ELIFA), and the like. In addition, immunological imaging methods capable of detecting dental caries are also provided by the invention, including but not limited to a colloidal-gold based colorimetric assay, and radioscintigraphic imaging methods using radiolabeled SWLA1, SWLA2, and SWLA3 antibodies (e.g., U.S. Patent No. 4,920,059 issued April 24, 1990; U.S. Patent No. 5,079,172 issued January 7, 1992). In addition the antibodies of the invention can be conjugated with other dyes or fluorescent markers and used

20

25

30



directly on the tooth to image caries. Such assays may be clinically useful in the detection and monitoring of dental caries. Such assays generally comprise using one or more of the SWLA1, SWLA2, and SWLA3 antibodies.

5 In addition to the immunological assays and imaging methods, the invention also includes an immunoconjugate comprising a molecule containing the antigen-binding region of the SWLA1, SWLA2, or SWLA3 antibody, or a fragment thereof containing the antigen binding region, joined to for example a therapeutic agent, a diagnostic agent or a cytotoxic agent for treatment of dental caries. Examples of  
10 cytotoxic agents include, but are not limited to, chlorhexidine, fluoride, ricin, doxorubicin, daunorubicin, taxol, ethidium bromide, mitomycin, etoposide, tenoposide, vincristine, vinblastine, colchicine, dihydroxyanthracenedione, actinomycin D, diphtheria toxin, *Pseudomonas* exotoxin (PE) A, PE40, abrin, glucocorticoid and radioisotopes.

15 The SWLA1, SWLA2, and SWLA3 monoclonal antibodies of the invention are useful for diagnostic applications, both *in vitro* and *in vivo*, for the detection of dental caries. *In vitro* diagnostic methods are well known in the art (see, e.g., Roth, *supra* 1986, and Kupchik, *supra* 1988), and include immunohistological  
20 detection of dental caries or serologic detection of *S. mutans* (e.g., in saliva samples or other biological fluids).

Immunohistological techniques involve contacting a biological specimen, such as a saliva, tartar, or plaque specimen, with the antibody of the  
25 invention and then detecting the presence in the specimen of the antibody complexed to its antigen. The formation of such antibody-antigen complexes with the specimen indicates the presence of the antigen, *S. mutans*. Detection of the antibody in the specimen can be accomplished using techniques known in the art, such as the immunoperoxidase staining technique, the avidin-biotin (ABC) technique or  
30 immunofluorescence techniques (Ciocca et al., (1986); Helstrom et al., (1986); and Kimball (ed.), (1986)).

Serologic diagnostic techniques involve the detection and quantitation of *S. mutans* antigens that have been secreted or "shed" into the saliva or other biological fluids of patients with dental caries. Such antigens can be detected in the saliva using techniques known in the art such as radioimmunoassays (RIA) or enzyme-linked immunosorbent assays (ELISA) wherein an antibody reactive with the "shed" antigen is used to detect the presence of the antigen in a fluid sample (see, e.g., Uotila et al., (1981) and Allum et al., 1986). These assays, using the antibodies disclosed herein, can therefore be used for the detection of *S. mutans* in biological fluids. Thus, it is apparent from the foregoing that the antibodies of the invention can be used in most assays involving antigen-antibody reactions. These assays include, but are not limited to, standard RIA techniques, both liquid and solid phase, as well as ELISA assays, immunofluorescence techniques, and other immunocytochemical assays (see, e.g., Sikora et al. (1984)).

15

The antibodies of the invention are also useful for *in vivo* diagnostic applications for the detection of dental caries. One such approach involves the detection of dental caries *in vivo* by imaging techniques using the antibody labeled with an appropriate imaging reagent that produces a detectable signal when bound to *S. mutans*. Imaging reagents and procedures for labeling antibodies with such reagents are well known (see, e.g., Wensel and Meares, (1983); Colcher et al., (1986)). The labeled antibody may be detected by a technique such as radionuclear scanning (see, e.g., Bradwell et al. (1985)).

20

The antibody fragments used in the immunoconjugates can include Fv, Fab, Fab' or F(ab)'<sub>2</sub> fragments. Use of immunologically reactive fragments, such as the Fv, Fab, Fab', or F(ab)'<sub>2</sub> fragments, is often preferable, especially in a therapeutic context, as these fragments are generally less immunogenic than the whole immunoglobulin. These antibodies, as well as unconjugated antibodies, may be useful therapeutic agents naturally targeted to *S. mutans* cells to kill the cells, thus preventing and or treating dental caries resulting from the accumulation of *S. mutans*. Techniques

25

30

for conjugating therapeutic agents to antibodies are well known (see, e.g., Arnon et al., 1985; Hellstrom et al. 1987; Thorpe, (1985); and Thorpe et al., (1982)).

The SWLA1, SWLA2, and SWLA3 antibodies may also be used in  
5 methods for purifying *S. mutans* proteins and peptides and for isolating homologues and related molecules. Methods for purification of proteins and peptides using antibodies as capture reagents are well known in the art. For example, in one embodiment, a method of purifying an *S. mutans* protein comprises incubating a  
SWLA1, SWLA2, or SWLA3 antibody, which has been coupled to a solid matrix, with  
10 a lysate or other solution containing *S. mutans* proteins or peptides, under conditions which permit the SWLA1, SWLA2, or SWLA3 antibody to bind to the *S. mutans* protein or peptides; washing the solid matrix to eliminate impurities; and eluting the *S. mutans* proteins or fragments from the coupled antibody.

15 The invention further includes a method for detecting the presence of *S. mutans* on teeth in a subject or in a saliva, plaque, or tartar sample from a subject, comprising contacting at least one tooth or the sample with the SWLA1, SWLA2, or SWLA3 antibody and detecting the binding of the antibody with the *S. mutans* on the tooth and or in the sample. The antibody can be administered by topical application to  
20 the surface of the teeth by means including in a toothpaste, mouthwash, lozenge, gel, powder, spray, liquid, tablet, or chewing gum. One can detect the presence of *S. mutans* by determining the presence of a complex formed between the monoclonal antibodies and *S. mutans* cells as a result of contacting the tooth and or the sample with a labeled antibody, the complex being indicative of the presence of *S. mutans* in the  
25 sample. The antibodies of the invention can be labeled so as to directly or indirectly produce a detectable signal. The label can for example be selected from the following compounds a radiolabel, an enzyme, a chromophore, a chemiluminescent moiety, a bioluminescent moiety, or a fluorescer. When a fluorescer is used the fluorescence can be detected by means of fluorescence microscopy, fluorometer, or by flow cytometry.  
30 A colloidal gold colorimetric system can also be used to detect the presence of *S.*

*mutans*. The colloidal gold system is well known in the art. (J.A.K. Hasan, et al. (1994); and E. Harlow, D. Lane. (1988)).

The invention also includes a method for diagnosing, in a subject, the  
5 early onset of dental caries. This can be accomplished by quantitatively determining on  
at least one tooth in a subject, or in a saliva, plaque, or tartar sample from a subject, the  
number of *S. mutans* present using an antibody of the invention and comparing the  
number of *S. mutans* cells so determined to the amount in a sample from a normal  
control, i.e. a subject free from dental caries. The normal range for *S. mutans* can be  
10 determined using any of the above detection methods (i.e. detecting labeled antibody to  
*S. mutans*) and quantifying the amount of *S. mutans* in a normal subject or subjects free  
of dental caries. For example, a normal range can be 1 cell/ml to approximately  $1 \times 10^5$   
cells/ml or  $1 \times 10^5$  cells/ml to  $1 \times 10^6$  cell/ml. Other ranges are possible. If the subject  
has a measurably higher amount of *S. mutans* present that is outside of the normal range  
15 it would indicate the early onset of dental caries in the subject.

The invention also includes a method for monitoring the course of dental  
caries in a subject. One can test teeth or a saliva, plaque, or tartar sample from a subject  
with the antibodies of the invention at different points in time and determine if there has  
20 been a change in the level of *S. mutans* present. An increase over a previous reading for  
that individual would suggest increased caries activity. For example if a first test of a  
subject's saliva sample gave a result of less than  $1 \times 10^5$  *S. mutans* cells/ml and a  
sample taken at a later time gave a result of greater than  $1 \times 10^5$  *S. mutans* cells/ml it  
would indicate that the subject now has an increased risk of dental caries.

25

The invention further comprises a method of protecting teeth from  
dental caries by topically applying an SWLA1, SWLA2, or SWLA3 antibody, or a  
fragment thereof containing the *S. mutans* antigen binding activity, to teeth of a subject.  
The antibody can be applied topically to the surface of the teeth by means of for  
30 example, of a toothpaste, mouthwash, lozenge, gel, powder, spray, liquid, tablet, or  
chewing gum formulated using standard methods. The antibody can be linked to a

toxic agent that kills the bacteria and applied to the surface of the teeth by, for example, any of the above methods. The proper dose of the monoclonal antibodies of the invention can be easily determined using methods which are well known to one skilled in the art (see, generally, Goodman et al. (ed.), 1993).

5

The methods described herein for detecting *S. mutans* may be performed using diagnostic kits (e.g., U.S. Patent No. 5,141,850 issued August 25, 1992; U.S. Patent No. 5,202,267 issued April 13, 1993; U.S. Patent No. 5,571,726 issued November 5, 1996; U.S. Patent No. 5,602,040 issued February 11, 1997). Such kits  
10 include at least one monoclonal antibody of the invention and reagents for detecting the binding of the monoclonal antibody to *S. mutans* cells present on teeth in or in a sample, e.g. of saliva, taken from a subject. The reagents include agents capable of detection, for example by fluorescence and ancillary agents such as buffering agents. The kits may also include an apparatus or container for conducting the methods of the  
15 invention and/or for transferring samples to a diagnostic laboratory for processing, as well as suitable instructions for carrying out the methods of the invention.

### ADVANTAGES OF THE INVENTION

20

With the monoclonal antibodies of the invention it is possible to monitor the detailed topology and proportion of *S. mutans* relative to other bacterial species during the course of plaque formation and the initiation and progression of carious lesions in a subject (e.g. with fluorescence microscopy). This in turn can lead to the  
25 development of improved treatment of dental caries. For example, antibodies of the invention can be conjugated with a regular or fluorescent dye. A solution containing such antibodies can be used to rinse a patient's mouth. The dyelinked antibodies can bind to the location of the dental caries. The dental caries image can be shown on a TV screen through a video or digital micro-camera.

30

With the fluorescent dye-linked monoclonal antibody and video imaging techniques, it is possible to label the bacteria at infection sites and thereby assist in detecting carious lesions at an early stage and in determining whether or not the lesion is active. This aids diagnosis, treatment and improves the management of dental health.

5

The monoclonal antibody based detection methods of the invention allows a rapid, accurate, and economic way to quantitatively measure the *S. mutans* in a subject, with significant advantages compared to current methods. As the first step towards development of effective and accurate caries risk assessment systems, we have described methods in this study that combine monoclonal antibodies with fluorometry techniques for detection and enumeration of *S. mutans*. These methods, especially flow cytometry, are able to rapidly detect the bacterium with high specificity and enumerate it with high accuracy. With these methods, it will be possible to process a large number of saliva samples in a short period of time at low cost. This will allow low cost, accurate assays to reevaluate the correlation between the salivary count of *S. mutans* and the presence and rate of progression of dental caries. Such assays can consists of monoclonal antibodies linked to a colloidal gold colorimetric system on test strips. The invention includes the use of a test system for rapid and simple assay of *S. mutans* by color change with simple immersion in fresh saliva. Such a method is suitable for use at a dentist's chairside as well as in the patient's household to assess dental caries risk. The accurate and objective assessment of dental caries risk state and/or caries activity state with any of these or with similar technologies will permit targeted preventive and curative treatment, thereby significantly improving human dental health.

25

In order that the invention described herein may be more fully understood, the following example is set forth. It should be understood that this example is for illustrative purposes only and is not to be construed as limiting the scope of this invention in any manner.

30

#### EXAMPLE 1

## MATERIALS AND METHODS

### Bacterial strains and culture conditions.

5 Bacterial strains used are listed in Table 2. *S. mutans* was grown in Brain-Heart Infusion (BHI) medium (Difco) with supplementations of haemin (5 µg/ml). Other bacteria were grown in various media as suggested by the American Type Culture Collection (ATCC). The anaerobic bacteria were grown in an atmosphere of 80% N<sub>2</sub>, 10% CO<sub>2</sub> and 10% H<sub>2</sub> at 37°C.

10

### Production and screening of species-specific monoclonal antibodies against *S. mutans*.

Type c *S. mutans* strain ATCC25175 were grown to log phase in BHI  
15 medium and washed twice with phosphate buffered saline pH 7.2 (PBS) by centrifugation at 3000 x g for 5 min. The pellet was resuspended in 1% formalin/0.9% NaCl, mixed at room temperature (RT) for 30 min and washed twice with 0.9% NaCl. BALB/c mice (8-10 weeks) were immunized intraperitoneally with 100 µl of the antigen containing approximately 10<sup>8</sup> whole cells of formalinized intact *S. mutans*  
20 bacteria emulsified with Freund's incomplete adjuvant (FIA). After 3-5 weeks they received a second dose of antigen (10<sup>8</sup> whole cells of bacteria in FIA). Three days prior to fusion, the mice were boosted intravenously with 10<sup>8</sup> whole cells in saline.

The standard tissue culture media was RPMI 1640 (Gibco) medium  
25 supplemented with 2 mM L-glutamine, 1 mM sodium pyruvate, and 10 mM HEPES and containing 100 µg/ml penicillin and 100 µg/ml streptomycin with 10% fetal calf serum. Hybrids were selected in media containing HAT (100 µg Hypoxanthine, 0.4 µM Aminopterin; 16 µM Thymidine). HT (100 µg Hypoxanthine, 16 µM Thymidine) was maintained in the culture medium for 2 weeks after aminopterin was withdrawn.  
30 OPI (1 mM oxaloacetate, 0.45 mM pyruvate and 0.2 U/ml bovine insulin) was added as additional growth factors to the tissue culture during cloning of hybridomas.

Hybridomas were raised according to the procedure reported by Kohler & Milstein (1975). The NSI/Ag4.1 mouse myeloma cell line was used as the fusion partner and grown in spinner cultures in 5% CO<sub>2</sub> at 37°C and maintained in log phase of growth prior to fusion.

5

The following approach was used for screening for species-specific monoclonal antibodies against *S. mutans*. The initial screening was performed using an ELISA assay, which selects for the culture supernatants containing antibodies that bind to *S. mutans*. Formalinized bacteria were diluted in PBS to OD<sub>600</sub>=0.5, and added to duplicate wells (100 µl) in 96 well PVC ELISA plates preincubated for 4 h with 100 µl of 0.02 mg/ml Poly-L-lysine. These antigen-coated plates were incubated overnight at 4°C in a moist box then washed 3 times with PBS and blocked with 0.5% fetal calf serum in PBS and stored at 4°C. A volume (100 µl) of mature hybridoma supernatants were added to the appropriate wells of the antigen plates, incubated for 1 hr at room temperature, washed 3 times with PBS-0.05% Tween 20, and bound antibody was detected by the addition of polyvalent goat-anti-mouse IgG antibody conjugated with alkaline phosphatase diluted 1:1000 with PBS-1% fetal calf serum. After the addition of the substrate, 1 mg/ml p-nitrophenyl phosphate in carbonate buffer (15 mM Na<sub>2</sub>CO<sub>3</sub>, 35 mM NaH<sub>2</sub>CO<sub>3</sub>, 10 mM MgCl<sub>2</sub> pH 9.6), the color development after 15 min was measured in an EIA reader at 405 nm. The positive supernatants (3 fold higher than control) were then subjected to the immunoprecipitation assay (mixing 100 µl bacteria with 100 µl supernatant) to screen for those with strong positive reactivity. These supernatants were then used to test crossreactivity with the following bacteria (strains listed in Table 2); *Streptococcus mutans*, *Streptococcus rattus*, *Streptococcus gordonii*, *Streptococcus mitis*, *Streptococcus sobrinus*, *Streptococcus sanguis*, *Streptococcus anginosus*, *Lactobacillus acidophilus*, *Lactobacillus casei*, *Actinobacillus actinomycetemcomitans*, *Porphyromonas gingivalis*, *Prevotella intermedia*, *Bacteroides forsythus*, *Eikenella corrodens*, *Fusobacterium nucleatum*, *Treponema denticola*, *Campylobacter rectus*, *Myxococcus xanthus*, and *Escherichia coli*. The results are shown in Table 2.

30



**Detecting *S. mutans* with fluorescence microscopy.**

A volume (10  $\mu$ l) of bacterial solution was mixed with 5  $\mu$ l of culture supernatant and incubated at room temperature for 20 seconds, then 1  $\mu$ l fluorescein isothiocyanate (FITC) linked goat-anti-mouse IgG antibody (Sigma) was added to the solution. The mixture was placed onto a Hausser bacterial counting chamber (Bright-line 1475) and observed with a fluorescent microscope (Zeiss Axiophot) for enumeration of the bacteria.

**10 Detecting *S. mutans* with a fluorometer.**

The bacteria were labeled with FITC molecules in the same way described above. The mixture was washed twice with PBS by centrifugation to remove the excess FITC-linked goat-anti-mouse antibody in solution. The pellet was then resuspended in PBS solution and put into a fluorometer (TD700, Turner Designs, Sunnyvale, CA) to measure the FITC fluorescent dyes bound to *S. mutans*, which reflects the bacterial concentration in the sample.

**Detecting *S. mutans* with flow cytometry.**

20

The bacteria were labeled with FITC molecules in the same way described above and analyzed with a Fluorescence-Activated Cell Sorter (FACS) (COULTER EPICS elite flow cytometer, Coulter Corp. Miami, FL). The FACS machine detects every particle in a solution and separates them based on their fluorescent intensity, with a capacity of 10,000 cells per second with an accuracy of 99.99%. With FITC-linked monoclonal antibodies against *S. mutans*, the bacteria can be easily detected and enumerated by the FACS machine in any bacterial mixture.

**30 Results**

**Isolation of species-specific monoclonal antibodies.**

Three BALB/c mice were immunized with formalinized *S. mutans* and used for production of monoclonal antibodies. The No. 2 mouse was selected because  
5 its serum showed the strongest positive reactivity with the bacterium and because 1835 mature hybridomas were obtained from this mouse.

Given the complexity of the surface of the *S. mutans* that contains a vast amount of distinct epitopes, it was important to have enough different hybridomas for  
10 binding assays in order to obtain some species-specific monoclonal antibodies. All 1835 mature hybridoma supernatants were screened with the ELISA assay, and 781 supernatants were found to have positive reactivity with *S. mutans*. Further immunoprecipitation assays identified 116 supernatants that gave the strongest positive reactivity. These culture supernatants were used to test cross-reactivity with the bacteria  
15 listed in Table 2.

-25-

TABLE 2

Species	Strain name	Cross-reactivity		
		SWLA1	SWLA2	SWLA3
<i>Streptococcus mutans</i>	ATCC25175	+	+	+
	LM7	+	+	+
	OMZ175	+	+	+
	ATCC31377	+	+	+
<i>Streptococcus rattus</i>	ATCC19645	-	-	-
<i>Streptococcus gordonii</i>	ATCC10558	-	-	-
	ATCC13396	-	-	-
<i>Streptococcus mitis</i>	ATCC49456	-	-	-
<i>Streptococcus sobrinus</i>	ATCC33478	-	-	-
	6715	-	-	-
<i>Streptococcus sanguis</i>	ATCC10556	-	-	-
	ATCC49295	-	-	-
<i>Streptococcus anginosus</i>	ATCC33397	-	-	-
<i>Lactobacillus acidophilus</i>	ATCC4356	-	-	-
<i>Lactobacillus casei</i>	ATCC4646	-	-	-
<i>A. actinomycetemcomitans</i>	ATCC33384	-	-	-
<i>Porphyromonas gingivalis</i>	ATCC33277	-	-	-
<i>Prevotella intermedia</i>	ATCC49046	-	-	-
<i>Bacteroides forsythus</i>	ATCC43037	-	-	-
<i>Eikenella corrodens</i>	ATCC23834	-	-	-
<i>Fusobacterium nucleatum</i>	ATCC25586	-	-	-
<i>Treponema denticola</i>	ATCC33520	-	-	-
<i>Campylobacter rectus</i>	ATCC33238	-	-	-
<i>Myxococcus xanthus</i>	DZ2	-	-	-
<i>Escherichia coli</i>	HB101	-	-	-

Three supernatants were identified which had the highest positive reactivity with *S. mutans*, yet did not have any significant cross-reactivity with other bacteria listed. The three hybridomas in the supernatants were further subcloned, purified, and designated as SWLA1, SWLA2, and SWLA3. Subclass isotype analysis indicates that all three monoclonal antibodies are of IgG subclass. Western blot analysis showed that the antibodies did not cross-react with any other bacterial surface proteins, distinguishing them from the known antibodies to *S. mutans* that cross-react with proteins on the cell surface of other bacteria. These findings indicate that the antibodies of the invention most likely bind to unique polysaccharide epitope on the surface of *S. mutans*.

#### Enumeration of *S. mutans* with fluorescence microscopy.

SWLA1, SWLA2 and SWLA3 antibodies were used to specifically label *S. mutans* in a mixture of bacterial cells. The monoclonal antibody-labeled *S. mutans* cells were then treated with FITC linked goat-anti-mouse IgG antibody to bind the monoclonal antibody and consequently label *S. mutans* cells with FITC. The FITC labeled *S. mutans* cells could be viewed and enumerated directly using a fluorescence microscope. Fig. 1(a). shows a microscopic image of a bacterial mixture containing *E. ratius*, *S. gordonii*, *S. mitis*, *S. sanguis*, *E. coli* as well as *S. mutans*. When the phase contrast lighting was shifted to fluorescent lighting, only *S. mutans* cells exhibited a fluorescent image due to bound FITC molecules, while other bacteria did not (Fig. 1(b).). In this way, the number of *S. mutans* in the mixture was easily recognized and enumerated. We compared this method of enumeration with the conventional colony counting method (the colony counting was performed by diluting bacteria with PBS and plating on BHI plates) and found good correlation between the two methods (Table 3).

-27-

TABLE 3

No. of bacteria assayed by colony counting <sup>1</sup> (cells/ml)	No. of bacteria assayed by fluorescent microscopy <sup>1</sup> (cells/ml)
1. $5.63 \times 10^6$	$4.8 \times 10^6$
2. $4.72 \times 10^5$	$5.2 \times 10^5$
3. $5.14 \times 10^4$	$5.0 \times 10^4$

<sup>1</sup>The numbers listed are the average of triplicates.

**Enumeration of *S. mutans* with fluorometer.**

As described above, *S. mutans* cells in a solution can be fluorescence-labeled with the monoclonal antibodies of the invention plus FITC-linked goat-anti-mouse IgG antibody, and the amount of fluorescence can be measured with a  
5 fluorometer. A linear correlation was observed between the amounts of fluorescence and concentrations of *S. mutans* in the solutions assayed by the conventional colony counting method (the colony counting was performed by diluting bacteria with PBS and plating on BHI plates.) (see Table 4).

-29-

TABLE 4

No. of bacteria assayed by colony counting <sup>1</sup> (cells/ml)	Fluorescence intensity detected by a fluorometer <sup>2,3</sup> (arbitrary unit)
1. $1.5 \times 10^6$	2262±12
2. $1.3 \times 10^5$	228±1
3. $1.6 \times 10^4$	23±1
4. $1.4 \times 10^3$	3±1

**Automated enumeration of *S. mutans* with fluorescence-activated cell sorter.**

The FITC labeled *S. mutans* cells were detected and enumerated with FACS. Figure 2 shows that the FITC-linked monoclonal antibodies which specifically  
5 bound *S. mutans* were effectively detected by FACS while other oral bacteria, such as *F. nucleatum* or *T. denticola* labeled with the same antibodies, showed no fluorescence, i.e. the antibodies did not bind to the other oral bacteria and were specific to *S. mutans*.

**Conclusion**

10

Applicants have discovered novel antibodies that are species-specific for *S. mutans* bacteria. These antibodies can be used to detect the presence of *S. mutans* in a subject thus allowing diagnose, treatment and preventative care for dental caries.

This invention includes methods for a quick and inexpensive test for the presence of *S.*  
15 *mutans*.



## References

Allum et al., Surg. Ann., (1986) 18:41-64.

Amon et al., "Monoclonal Antibodies For Immunotargeting Of Drugs In Cancer Therapy," Monoclonal Antibodies And Cancer Therapy, Reisfeld et al. (eds.), pp. 243-56 (Alan R. Liss, Inc. 1985).

Borrebaeck et al. *Proc. Nat'l. Acad. Sci. (USA)* (1988) 85:3995-99.

Bradwell et al., Monoclonal Antibodies for Cancer Detection and Therapy, Baldwin et al. (eds.), pp. 65-85, Academic Press (1985).

Brady, L. J. et al., "Identification of monoclonal antibody-binding domains within antigen P1 of *Streptococcus mutans* and cross-reactivity with related surface antigens of oral streptococci," *Infect. Immun.* 59:4425-4435 (1991).

Brown et al., *J Immunol.*, 127(2):539-46 (1981).

Brown et al., *J Biol. Chem.*, 255:4980-83 (1980).

Cancer Diagnosis In Vitro Using Monoclonal Antibodies, Kupchik (ed.) Marcel Dekker, Inc., New York (1988).

Carlen, A. and J. Olsson, "Monoclonal antibodies against a high-molecular-weight agglutinin block adherence to experimental pellicles on hydroxyapatite and aggregation of *Streptococcus mutans*," *J Dent. Res.* 74:1040-1047 (1995).

Cedars-Sinai Medical Center, Guide to laboratory services, Lexi-Comp Inc., Hudson, Ohio. (1995).

Chia, J. S., Lin, R. H., Lin, S. W., Chen, J. Y., Yang, C. S., "Inhibition of glucosyltransferase activities of *Streptococcus mutans* by a monoclonal antibody to a subsequence peptide," *Infect. Immun.* 61:4689-4695 (1993).

Ciocca et al., *Meth. Enzymol.* 121:562-79 (1986).

Clark, J. K., "On the bacterial factor in the etiology of dental caries," *Brit. J. Path.* 5:141-147 (1924).

Colcher et al., *Meth. Enzymol.*, 121:802-16 (1986).

Ellen, R.P., "Microbiological assays for dental caries and periodontal disease susceptibility," *Oral Sci. Rev.* 8:3-23 (1976).

Fink, et al. *Prog. Clin. Pathol.* 9:121-33 (1984).

Fitzgerald, R. J. and P. H. Keyes, "Demonstration of the etiologic role of streptococci in experimental caries in the hamster," *JADA* 67:9-19 (1960).

Fukushima K., Okada T., Ochiai K., "Production, characterization, and application to monoclonal antibodies which distinguish three glucosyltransferases from *Streptococcus mutans*," *Infect. Immun.*, 61:323-328 (1993).

Goodman and Gilman's The Pharmacological Basis of Therapeutics, 8<sup>th</sup> ed. (A. Gilman, T. Rall, A. Nies and P. Taylor: eds.) McGraw-Hill, Inc., 53:1272-1273 (1993).

Golub, E. S. and D. R. Green, *Immunology: A Synthesis*, Sinauer Associates, Inc. Sunderland, Massachusetts, 164-166 (1991).

Harlow E. and D. Lane, "Antibodies, a laboratory manual," CSH Press, Cold Spring Harbor Laboratory, New York (1988).

Hasan, J.A.K., A. Huq, M.L. Tamplan, R.J. Siebeling, and R.R. Colwell, "A novel kit for rapid detection of *Vibrio cholerae* O1," *J Clin. Microbiol.*, 32:249-252 (1994).

Hellstrom et al., *Cancer Research*, 46:3917-23 (1986).

Hellstrom et al., "Antibodies For Drug Delivery", in Controlled Drug, Delivery (2nd Ed.), Robinson et al. (eds.), pp. 623-53 (Marcel Dekker, Inc. 1987).

Hume, W. R., "Need for change in standards of caries diagnosis-perspective based on the structure and behavior of the carious lesion," *J Dent. Educ.* 57:439-443 (1993).

Igarashi T., Yamamoto A., Goto N., "Direct detection of *Streptococcus mutans* in human dental plaque by polymerase chain reaction," *Oral Microbiology and Immunology*, 11:294-298 (1996).

Kimball (ed.), Introduction To Immunology (2nd Ed.), pp. 113 -117, Macmillan Publ. Co (1986).

Kohler G., and C. Milstein, "Continuous cultures of fused cells secreting antibody of predefined specificity," *Nature*, 256:495-497 (1975).

Krasse, B., "The cariogenic potential of foods--a critical review of current methods," *Inter. Dent. J.* 35:36-42 (1985).

Krasse B., "Biological factors as indicators of future caries," *Inter. Dent. J.* 38:219-225 (1985).

Liu et al., *Proc. Natl. Acad. Sci. (USA)* 84:3439-43 (1987).

Loesche, W.J., "Dental Caries: a treatable infection," Thomas, Springfield (1982).

Loesche, W.J., "Role of streptococcus mutans in human dental decay," *Microbiol. Rev.* 50:353-380 (1986).

Loesche, W.J., Rowan, J., Straffon, L.H., Loos, P.J., "Association of *Streptococcus mutans* with human dental decay," *Infect. Immun.* 11: 1252-1260 (1975).

Monoclonal Antibodies in Cancer: Advances for Diagnosis and Treatment, Roth (ed.), Futura Publishing, Mt. Kisco, New York (1986).

Mundorff, S. A., Billings, R. J., Leverett, D. H., Featherstone, J. D., Gwinner, L. M., Shields, C. P., Proskin, H. M., Shaffer, C. L., "Saliva and dental caries risk assessment," *Ann. New York Acad Sci.* 694:302-304 (1993).

Oi et al., *Biotechnologies*, 4(3):214-221 (1986).

Sikora et al. (eds.), Monoclonal Antibodies, pp. 32-52, Blackwell Scientific Publications, (1984).

Shawler et al., *J Immunol.* 135:1530-35 (1985).

Tanzer, J. M., "Understanding dental caries: an infectious disease, not a lesion," *Inter. J Oral Biol.* 22:205-214 (1997).

Thorpe "Antibody Carriers Of Cytotoxic Agents In Cancer Therapy: A Review", in Monoclonal Antibodies '84: Biological And Clinical Applications, Pinchera et al. (eds.), pp. 475-506 (1985).

Thorpe et al., "The Preparation And Cytotoxic Properties Of Antibody-Toxin Conjugates", Immunol. Rev., 62:119-58 (1982).

Uotila et al., *J Immunol. Methods*, 42:11 (1981).

Van Houte, J., "Microbiological predictors of caries risk," *Adv. Dent. Res.* 7:80-86 (1993).

Wensel and Meares, Radio Immunoimaging and Radioimmunotherapy, Esvier, New York (1983).


Yeh et al., *Proc. Nat'l. Acad Sci. (USA)*, 76 (6):2927-31 (1976).

Yeh et al., *Int. J Cancer*, 29:269-75 (1982).


Zola et al., "Monoclonal Hybridoma Antibodies: Techniques And Applications", Hurell (ed.) pp. 51-52 (CRC Press 1982).

What is claimed is:

1. A monoclonal antibody that specifically binds an antigen on the surface of *Streptococcus mutans* and which is the monoclonal antibody produced by a hybridoma deposited with the American Type Culture Collection as ATCC No. HB12559, and which is designated SWLA1.
2. A monoclonal antibody that specifically binds an antigen on the surface of *S. mutans* and which is the monoclonal antibody produced by a hybridoma deposited with the American Type Culture Collection as ATCC No. HB 12560, and which is designated SWLA2.
3. A monoclonal antibody that specifically binds an antigen on the surface of *S. mutans* and which is the monoclonal antibody produced by a hybridoma deposited with the American Type Culture Collection as ATCC No. HB 12258, and which is designated SWLA3.



*FIG. 1A*



*FIG. 1B*

2/2

FIG. 2a




FIG. 2b




FIG. 2c




FIG. 2d




FIG. 2e





FIG. 2f





## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/19034

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 C07K16/12

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	W. SHI ET AL.: "Rapid and quantitative detection of Streptococcus mutans with species-specific monoclonal antibodies." HYBRIDOMA, vol. 17, no. 4, August 1998 (1998-08), pages 365-371, XP002119822 New York, NY, USA the whole document	1-3
X	J. DE SOET ET AL.: "Identification of mutans streptococci with monoclonal antibodies." ANTONIE VAN LEEUWENHOEK, vol. 58, no. 4, November 1990 (1990-11), pages 219-225, XP002119823 Amsterdam, The Netherlands the whole document	1-3



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

## \* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

21 October 1999

Date of mailing of the international search report

05/11/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Nooij, F

## INTERNATIONAL SEARCH REPORT

Interr      al Application No

PCT/US 99/19034

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	J. DE SOET ET AL.: "Streptococcus sobrinus and Streptococcus mutans in a longitudinal study of dental caries." MICROBIAL ECOLOGY IN HEALTH AND DISEASE, vol. 6, no. 5, September 1993 (1993-09), pages 237-243, XP002119824 Chichester, GB the whole document ----	1-3
X	Y. AKIYAMA ET AL.: "Monoclonal antibodies specific for a polysaccharide antigenic determinant of Streptococcus mutans." KYUSHU SHIKA GAKKAI ZASSHI, vol. 49, no. 5, 1995, pages 360-372, XP002119825 Japan abstract ----	1-3
X	DATABASE WPI Week 9033 Derwent Publications Ltd., London, GB; AN 1990-252062 XP002119826 & JP 02 177898 A (NAGASE SANGYO KK), 10 July 1990 (1990-07-10) abstract ----	1-3
X	DATABASE WPI Week 9033 Derwent Publications Ltd., London, GB; AN 1990-252063 XP002119827 & JP 02 177899 A (NAGASE SANGYO KK), 10 July 1990 (1990-07-10) abstract ----	1-3
X	DATABASE WPI Week 9816 Derwent Publications Ltd., London, GB; AN 1998-174917 XP002119828 & JP 10 036400 A (LION CORP.), 10 February 1998 (1998-02-10) abstract -----	1-3

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/19034

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
JP 2177898 A	10-07-1990	JP 2749338 B	13-05-1998
JP 2177899 A	10-07-1990	NONE	
JP 10036400 A	10-02-1998	NONE	

Form PCT/ISA/210 (patent family annex) (July 1992)





*FIG. 1A*



*FIG. 1B*

2/2

FIG. 2a




FIG. 2b




FIG. 2c




FIG. 2d




FIG. 2e




FIG. 2f

